




Diabetes In LANCASTER County, 2005



Produced by the South Carolina Department of Health and Environmental Control

Impact of Diabetes in the United States


- Diabetes affects more than 23 million Americans and contributes to almost 234,000 deaths a year.
- Diabetes is becoming more common in the United States. From 1980 through 2007, the number of Americans with diabetes increased from 5.6 million to 17.9 million
- The direct and indirect costs of diabetes in the U.S. were nearly \$174 billion in 2007.
- 72,507 people died of diabetes in the U.S. in 2006.

Diabetes in South Carolina

- Approximately 332,000 adults in South Carolinians are diagnosed with diabetes. African Americans have twice the prevalence than Whites.
- The total direct costs of hospitalizations and emergency room visits were over \$8.5 billion in 2005.
- Diabetes is the seventh leading cause of death in South Carolina, claiming 1,183 lives in 2005.


Behavioral Risk Factors

- Major behavioral risk factors, such as overweight, physical inactivity and unhealthy diet, are partially responsible for development of type 2 diabetes (Fig 1).



Diabetes Prevalence

- An estimated 5099 adults (10.6% of adults) in Lancaster County suffer annually from Diabetes (Fig 2).




Hospital and Emergency Room Discharges

- In 2005, there were 130 hospitalizations for Diabetes in Lancaster County for all ages.
- There were an additional 1246 hospital discharges with diabetes-related conditions in Lancaster County.

- In 2005, there were 41 emergency room (ER) visits for diabetes as the primary diagnosis, among which 28 (68%) were by African-American patients.

- There were 204 ER visits for diabetes as a related condition.




- Medicare paid for 69% of the ER and Hospitalization charges in Lancaster County (Fig 3).

Complications

- Almost all patients (97%) hospitalized in SC for diabetes had complications in 2005; 26% of hospitalizations in Lancaster County (Table 1).
- In 2005, nearly three-quarters of people in SC with diabetes had hypertension, compared with only one-fourth of people without diabetes.


Table 1. Hospitalizations for Diabetes Complications in Lancaster County in 2005.			
Diabetes Complication	Hospital Admissions	Percent of Diabetes Hospital Admissions	Total costs
Renal Failure	227	16%	\$8,344,800
Lower Extremity Amputation	17	1%	\$1,023,100
Keto-acidosis	25	2%	\$708,100
Retinopathy	88	6%	\$3,000,800

- In 2005, more than 60% of people with diabetes have high cholesterol.
- In 2005, the rates for lower extremity amputations among blacks in SC were four times the rates among whites.




Mortality

- Diabetes is the eleventh leading cause of death in Lancaster County.
- A total of 9 people in Lancaster County died from diabetes in 2005.



- The Lancaster County mortality rate was 13.8/100,000 in 2005, below the state average.
- Lancaster County ranks #45 in diabetes mortality in South Carolina 2003-2005.



For More Information about diabetes contact:

Division of Diabetes Prevention and Control:

DHEC Bureau of Community Health and Chronic Disease Prevention

<http://www.dhec.sc.gov/health/chcdp/diabetes/index.htm>

(803) 545-4471

Diabetes Initiative of South Carolina

<http://www.musc.edu/diabetes/>

Generated by: Office of Chronic Disease Epidemiology,
Bureau of Community Health and Chronic Disease
Prevention

http://www.scdhec.gov/hs/epidata/state_reports.htm



Data Sources:

For more details, please view the Technical Document below

www.scdhec.gov/health/epidata/docs/EpiTechNotes.pdf

Hospital Discharges and ER Visits, contact:

Office of Research and Statistics SC
Budget and Control Board
(803) 898-9958

